

Summary

The purpose of this manuscript is to review and summarize the results of experiments conducted in our laboratory regarding the mechanism of meat tenderization during postmortem storage of carcasses at refrigerated temperatures. Clearly, the conversion of muscle to meat and the subsequent tenderization process are complex phenomena and much remains to be learned. However, current experimental data suggest that proteolysis of key myofibrillar proteins is the principal reason for improvement in meat tenderness during postmortem storage. Speculatively, the weakening and/or degradation of Z-disks and degradation of desmin (and probably degradation of titin) are responsible for the increased fragility of myofibrils during postmortem storage. There is substantial experimental evidence suggesting that the calpain proteolytic system is responsible for postmortem proteolysis that results in meat tenderization. Calpain is the only proteolytic system that has all of the characteristics that are necessary for bringing about postmortem changes that result in meat tenderization. Undoubtedly, other factors (such as rate of pH and temperature decline during rigor development, ionic strength, and others) influence the process. However, we believe that the rate and extent of postmortem proteolysis best explain the observed variation in tenderness at a constant age. Therefore, research efforts should be directed toward understanding the regulation of the calpain proteolytic system in postmortem muscle.

Introduction

The improvement in meat tenderness during postmortem storage of carcasses at refrigerated temperatures has been known since the turn of the century (LEHMANN, 1907). However, the mechanism through which these changes are brought about has remained elusive and controversial.

Because consumers consider tenderness to be the most important organoleptic characteristic of meat, it is essential that we understand the mechanism of meat tenderization so that methodologies can be developed to manipulate the process advantageously. Undoubtedly, the mechanism of tenderization is complex and affected by a number of variables. Over the years, the following variables have been proposed to influence meat tenderness: animal age and gender, rate of glycolysis, amount and solubility of collagen, sarcomere length, ionic strength and degradation of myofibrillar proteins.

The purpose of this manuscript is to review and summarize the results of experiments conducted in this laboratory related to the role of endogenous proteinases in the postmortem tenderization process. The manuscript is not intended to be a comprehensive review of all factors affecting meat tenderness. Throughout this manuscript, postmortem storage is defined as holding of carcasses at refrigerated temperatures and should be distinguished from other methods such as high temperature conditioning. Also, our research efforts have been directed toward understanding the causes of variation in meat tenderness of animals slaughtered at similar ages and should not be extrapolated to other situations. For additional information, the reader is referred to a number of review papers written on this subject (ASGHAR and BHATTI, 1987; DAVEY, 1983; DUTSON, 1983; DUTSON and PEARSON, 1985; GOLL et al., 1983; GREASER, 1986; KOOHMARAIE, 1988, 1992a,b; MARSH, 1977, 1983;

MARSH et al., 1988; OUALI, 1990, 1992; PEARSON, 1986; PENNY, 1980; ROBSON and HUIATT, 1983; ROBSON et al., 1981, 1984). Throughout this manuscript, due to space limitation, original information source will be given only when the subject has not been addressed in these review articles.

Postmortem Changes in Skeletal Muscle

Because of the number of recent reviews in this area (see above), only important changes as they relate to the objective of this manuscript will be discussed. During postmortem storage of carcasses, numerous changes occur in skeletal muscle, some of which result in the loss of tissue integrity which is translated into the improvement of meat tenderness. These changes include: 1) Z-disk weakening and/or degradation which leads to fragmentation of myofibrils. 2) Degradation of desmin which leads to fragmentation of myofibrils, probably through disruption of transverse crosslinking between myofibrils. 3) Degradation of titin. Titin filaments, which are made up of titin molecules, connect myosin filaments, along their length, from the M-line to the Z-disk (WANG, 1985). Titin has been proposed to be involved in the regulation of the elasticity of the muscle (WANG et al., 1991). When titin was preferentially destroyed by radiation (HOROWITS et al., 1986) or controlled proteolysis (YOSHIOKA et al., 1986), the tension of stretched muscle was reduced. Therefore, degradation of titin during postmortem storage would cause weakening of myofibril strength and, therefore, improvement in meat tenderness. 4. Degradation of nebulin. Because of the location of nebulin in myofibrils (I-band), it is not clear how nebulin degradation will affect meat tenderness. 5) Disappearance of troponin-T and simultaneous appearance of polypeptides with molecular weight of 28 to 32. This is the most noticeable and reported change that occurs during postmortem storage. However, because of the location of troponin-T in myofibrils (i.e., I-band), it is doubtful that degradation of troponin-T by itself will have a direct effect on meat tenderness. But, these changes (i.e., the disappearance of troponin-T and appearance of 28 to 32 kDa polypeptides) seem to be good indicators of the extent of postmortem proteolysis. The origin of 28 to 32 kDa polypeptides has not been determined and, therefore, these polypeptide could be from degradation of any myofibrillar proteins with molecular mass greater than 32 kDa. 6) Appearance of a polypeptide with a molecular weight of 95. Neither the origin nor its significance to meat tenderness is known. 7) Perhaps the most important observation is that the major contractile proteins (myosin and actin) are not affected. One of the important changes that occurs in the tissue is the ease of fragmentation of myofibrils under controlled homogenization, which does not occur in the unaged tissue. This phenomenon, first reported by DAVEY and GILBERT (1969), which is measured routinely by a number of laboratories, is called Myofibril Fragmentation Index (MFI) and is highly related to meat tenderness (for review see PARRISH, 1977). Speculatively, the weakening and/or degradation of Z-disks and degradation of desmin (and probably degradation of titin) are responsible for the increased fragility of myofibrils during postmortem storage.

Mechanisms of Postmortem Changes in Muscle Tissue

Clearly, the changes discussed in the previous section are all produced by proteolytic action; and, therefore, the changes resulting in improvement in meat tenderness are produced by endogenous proteases. This is not a new concept. As early as 1917, HOAGLAND et al concluded that proteolysis was an important factor contributing to postmortem changes in skeletal muscle, including meat tenderness. Because the proteolytic

changes that occur in skeletal muscle during postmortem storage are minimal, the classical methods failed to detect these changes and, therefore, the proteolysis hypothesis was questionable until the advent of gel electrophoresis. Gel electrophoresis made it possible to demonstrate these minimal, yet significant, changes and, therefore, give credibility to the proteolysis theory. Based on the observation reported by numerous laboratories, PENNY (1980) concluded that, "there is no doubt that proteolytic enzymes are responsible for the changes during conditioning (postmortem storage)." Tenderness could also be improved by changes in the connective tissue; however, because proteolytic changes in collagen (the principal component of the connective tissue) during postmortem storage comparable to those of myofibrillar proteins have not been observed (TARRANT, 1987), the role of collagen is questionable at best. In addition, while collagen may affect meat tenderness of the same muscle obtained from young (e.g., 1 year old) and old animals (e.g., 7 years old), it is doubtful if any significant differences exist in collagen solubility of muscle (e.g., longissimus) from animals of similar age. We, therefore, have concluded that differences in the rates of myofibrillar protein degradation are the principal reason for the observed variation in tenderness of meat obtained from animals of similar age.

Indeed, there is substantial experimental evidence in support of this theory. Some of these include: 1) Infusion of carcasses with zinc chloride, which inhibits postmortem proteolysis, also inhibits the tenderization process (KOOHMARAIE, 1990). 2) Muscle from β -adrenergic agonist fed animals which undergo minimal or no postmortem proteolysis, also remains tough compared to muscle from untreated animals (FIEMS et al., 1990; KRETCHMAR et al., 1990; KOOHMARAIE et al., 1991a; KOOHMARAIE and SHACKELFORD, 1991; WHEELER and KOOHMARAIE, 1992). 3) Differences in the extent of postmortem proteolysis are probably the reason for differences in meat tenderness between Bos taurus and Bos indicus breeds of cattle (SHACKELFORD et al., 1991; WHIPPLE et al., 1990) and 4) differences in rate of postmortem proteolysis are probably the reason for the observed differences in meat tenderness from pigs, sheep and cattle (KOOHMARAIE et al., 1991b).

Proteinases Involved in Postmortem Proteolysis

Currently, while the proteolysis theory is accepted by most, the question of proteinases involved has remained controversial. Proteinases should have the following characteristics to be considered as possible candidates for bringing about postmortem changes that result in meat tenderization: 1) be located within the skeletal muscle cell (for details see GOLL et al., 1983); 2) have access to the substrate (i.e., myofibrils); and 3) have the ability to degrade the same proteins that are degraded during postmortem storage. The proteolytic systems that have the potential to be involved in postmortem proteolysis include: 1) the lysosomal cathepsins; 2) the multicatalytic proteinase complex (MCP); and 3) the calpains. Current experimental evidence suggests that lysosomal cathepsins do not play a significant role in postmortem proteolysis. Some of this experimental evidence includes: 1) postmortem storage has no effect on actin and myosin, while among myofibrillar proteins these are the primary substrates for lysosomal cathepsins; 2) lysosomal cathepsins are normally located within lysosomes and must, therefore, be released to have access to myofibrils. While it has been assumed that during postmortem storage lysosomes are ruptured, thereby releasing cathepsins into the cytosol, there is no experimental evidence to support this hypothesis. To the contrary, the only experiment that has examined the accuracy of this hypothesis indicates that even after electrical stimulation and 28 days of storage at 4°C, lysosomal enzymes were still localized within lysosomes (LaCOURT et al., 1986). Because of

these and other reasons (KOOHMARAIE 1988, 1990, 1992a), we have concluded that lysosomal cathepsins do not play a significant role in postmortem proteolysis.

The second candidate is the MCP. Until recently, no experimental data were available to determine the role of MCP in this process. We have recently purified and characterized MCP from ovine skeletal muscle (KOOHMARAIE, 1992d). Our results indicate that ovine skeletal muscle indeed contains MCP with similar biochemical properties to MCP isolated from other mammalian and non-mammalian tissues. Some of the characteristics of ovine skeletal muscle MCP include: 1) molecular mass of 600 kDa which dissociates into a series of low molecular polypeptides ranging from 21 to 31 kDa; 2) it has no proteolytic activity as isolated from tissue, but it can be reversibly activated by heating at 60°C and with pre-treatment with low concentration of sodium dodecyl sulfate (SDS); 3) maximum proteolytic activity is observed at pH 7.5 to 8.0 and 45°C, it retains about 2% of its maximum activity at 5°C and pH 7.5, and about 22% of its maximum activity at pH 5.5 and 45°C; and 4) calcium chloride has no effect on its proteolytic activity. For more details of MCP characteristics, the reader is referred to excellent reviews (RIVETT, 1989; ORLOWSKI, 1990). Results of our experiments indicate that even after activation (by heating or incubation in the presence of SDS), myofibrils were very poor substrate for MCP. We incubated myofibrils with MCP and analyzed the effects with SDS-PAGE, phase and electron microscopy. Morphologically, MCP had no effect on myofibrils and based on SDS-PAGE data, MCP only degraded troponin-C and myosin light chain-1 and -2. These results indicated that MCP does not play a major role in postmortem proteolysis that results in meat tenderization.

In contrast to lysosomal proteinases and MCP, substantial experimental evidence exists suggesting that calpains are the primary proteolytic system responsible for postmortem proteolysis that results in meat tenderization. There is considerable experimental evidence indicating that calcium causes weakening and(or) degradation of Z-disks. The first report that documented the role of calcium in Z-disk weakening was that of DAVEY and GILBERT (1969). They reported that EDTA inhibited the weakening and disappearance of Z-disks and speculated that EDTA probably acts by chelating calcium. BUSCH et al (1972) provided further support by demonstrating that myofibril fragmentation was inhibited by EDTA and was induced by calcium. KOOHMARAIE et al (1988a) also demonstrated that all postmortem changes were completed within 24 hours when muscle slices were incubated with a buffer solution containing calcium chloride and none of the postmortem changes occurred if EDTA was included in the buffer instead of calcium chloride.

Acceleration of Postmortem Proteolysis and Tenderization Processes

Based on the observation reported in the previous section, it became evident that the elevation of calcium concentration in postmortem muscle is the cause of postmortem tenderization. To determine whether these observations could be repeated in situ, lamb carcasses were infused with a solution of calcium chloride to increase intracellular concentration of calcium (KOOHMARAIE et al., 1988b). Results indicated that postmortem proteolysis and tenderization were accelerated such that ultimate tenderness values were obtained within 24 hours of postmortem storage as opposed to 7 to 14 days in non-infused carcasses. Though these experiments were designed to activate calpains (KOOHMARAIE et al., 1988a,b, for review see KOOHMARAIE, 1988, 1992a), we do not know the precise mechanism(s) through which calcium chloride infusion accelerates postmortem proteolysis and

tenderness. However, we believe the primary mode of action of calcium is through activation of calpains (for review see KOOHMARAIE, 1988, 1992a). There is no doubt that calcium will induce changes other than activation of calpains, however, these changes may not affect meat tenderness (TAYLOR and ETHERINGTON, 1991; WHIPPLE and KOOHMARAIE, 1992). Regardless of the mechanism of action, calcium chloride infusion of whole carcasses or injection of cuts of meat is a very effective method of rapidly producing uniformly tender meat. The process has been very effective under all experimental conditions thus far examined, including: lamb carcasses (KOOHMARAIE et al., 1988b, 1989; ST. ANGELO et al., 1991); *Bos indicus* carcasses (KOOHMARAIE et al., 1990); beef round muscles (WHEELER et al., 1991); and mature cow carcasses (MORGAN et al., 1991); postrigor injection of beef longissimus muscle (WHEELER et al., 1992) and postrigor marination of beef steaks (WHIPPLE and KOOHMARAIE, 1992).

Current experimental data suggest that of the three proteolytic systems discussed, the calpain proteolytic system is the best possible candidate for causing postmortem proteolysis and tenderization because:

- 1) calpains have an absolute requirement for calcium and, clearly, the elevation of calcium is the reason for the observed changes in postmortem muscle that result in tenderization;
- 2) calcium has no effect on the activity of MCP (KOOHMARAIE, 1992d);
- 3) calcium not only does not stimulate cathepsins activity, but at 10 mM inhibits cathepsin B activity by 39% (BARRETT, 1973);
- 4) of these three proteolytic systems, the calpains are the only system that precisely reproduces postmortem changes under *in vitro* conditions; and
- 5) of these three proteolytic systems, calpain and MCP are localized in the cytosol, but cathepsins are located within lysosomes. While precise location of MCP in relation to myofibrils is not known, calpains are localized primarily at the Z-disk (for μ -calpain: 66% on Z-disk, 20% in I-band, and 14% in A-band; KUMAMOTO et al., 1992).

Conclusions

Clearly, the process of conversion of muscle to meat and the subsequent tenderization process is complex and much remains to be learned. Over the years, a number of factors have been proposed to influence the ultimate meat tenderness. These include rate of glycolysis, pH, sarcomere length, amount and solubility of collagen and postmortem proteolysis. Undoubtedly, all of these parameters and their interaction need to be considered to explain the observed variation in meat tenderness. Based on our current knowledge of the process, postmortem proteolysis is the most important of all and that most other factors (such as rate of glycolysis, ultimate pH, rate of temperature decline) affect meat tenderness by their influence on the proteolytic systems involved. Factors such as ionic strength and collagen solubility are probably involved in the process, but they cannot explain the differences observed in tenderness of meat obtained from animals of similar age. Rather, these factors set the so-called "background toughness" (MARSH, 1977). For any theory to be valid, it must be able to explain the large variation observed in tenderness of meat from animals of identical backgrounds (e.g., variation observed in tenderness of meat from *Bos Taurus* cattle slaughtered at 16 to 18 month of age). Clearly, collagen cannot explain these differences and neither can ionic strength. While ionic strength of postmortem muscle is double that of living tissue (equivalent of 165 and 280 mM NaCl; WINGER and POPE, 1980-81) and that such a significant elevation in ionic strength would be expected to affect myofibrils which may lead to their instability, the question that needs to be addressed is: why would ionic

strength be different in longissimus of animals of identical backgrounds. Let's examine two cases to see which of these three factors (collagen, ionic strength, and postmortem proteolysis) can explain the variation in meat tenderness.

Case 1: Tenderness of meat from β -adrenergic agonist-fed (BAA; L644,969 from Merck Sharp and Dohme) animals. When lambs (KRETCHMAR et al., 1990; KOOHMARAIE and SHACKELFORD, 1991; KOOHMARAIE et al., 1991a) and steers (WHEELER and KOOHMARAIE, 1992) are fed BAA, the meat from their carcasses is tough and postmortem storage has no effect on it (i.e., it remains tough). Ultimate pH is proposed to cause elevation of ionic strength in postmortem muscle. The correlation between ionic strength and pH is reported to be about -0.90 (for review see OULAI, 1990). Since BAA feeding does not affect ultimate pH of the muscle, one would expect that BAA should not effect ionic strength, yet meat from BAA fed animals is not affected by postmortem storage (i.e., remains tough). Because the half-life of collagen is in excess of 200 days and BAA effects are evident after 2 weeks of feeding (PRINGLE et al., 1992), toughness of meat from BAA-fed animals cannot be due to changes in collagen and, indeed, our data support this speculation (KOOHMARAIE and SHACKELFORD, 1991). However, all data collected thus far indicate that lack of postmortem proteolysis is the reason for the toughness of meat from BAA fed animals (FIEMS et al., 1990; KOOHMARAIE et al., 1991a; KOOHMARAIE and SHACKELFORD, 1991; KRETCHMAR et al., 1990; WHEELER and KOOHMARAIE, 1992).

Case 2: Toughness of meat from Bos indicus as compared to meat from Bos taurus. It has clearly been documented that meat obtained from Bos indicus carcasses is significantly tougher than that obtained from Bos taurus carcasses (RAMSEY et al., 1961; KOCH et al., 1982; PEACOCK et al., 1982; CROUSE et al., 1987, 1989). To identify the cause of these differences in tenderness, we (SHACKELFORD et al., 1991; WHIPPLE et al., 1990) determined a number of factors that are proposed to affect tenderness in meat obtained from Bos taurus and Bos indicus cattle raised under identical management practices (similar climate, diet, and slaughter age). Of all factors examined (pH and temperature decline, muscle composition, fiber type composition and diameter, amount and solubility of collagen, sarcomere length, MFI and SDS-PAGE of myofibrillar proteins during postmortem storage), only postmortem proteolysis, determined by MFI and SDS-PAGE, was different. Because neither the pattern of pH decline nor ultimate pH was different, it was concluded that ionic strength (see above discussion on the relationship between pH and ionic strength) is not the cause of differences in tenderness of meat from these breeds of cattle. Data clearly suggest that the reduced rate of postmortem proteolysis in meat from Bos indicus carcasses was the only logical explanation for differences in tenderness of meat from these two breeds of cattle (WHEELER et al., 1990; WHIPPLE et al., 1990).

Clearly, these two examples indicate that differences in the rate of postmortem proteolysis is the best explanation for the observed variation in meat tenderness.

Current experimental data suggest that the calpain proteolytic system is probably responsible for postmortem changes that result in improvement in meat tenderness. To manipulate the process, we must understand how calpains are regulated in postmortem muscle. Using a modeling approach, DRANSFIELD (1992) demonstrated that 68% of the variation in toughness was accounted for by variation in μ -calpain activity. Identification of the regulatory mechanism for calpain in postmortem muscle could enable us to manipulate the process and, thereby, enhance the tenderization process. Recently, we have begun to determine the regulation

of calpain in postmortem muscle (KOOHMARAIE, 1992c). Results indicate that pH and temperature, two key changes that occur in muscle during rigor development, have a dramatic effect on the inactivation of μ -calpain. We hope that such experimental approaches would lead to development of alternative carcass handling procedures during slaughter and early postmortem to maximize calpain potential and, therefore, improvement in the rate of tenderization.

Finally, we must develop the methodology to predict meat tenderness as early postmortem as possible and, ultimately, prior to slaughter. The development of such methodology would enable us to decide how a particular carcass should be marketed, depending upon its predicted eating quality. Variation in meat tenderness at the consumer level is one of the biggest problems that our industry is facing now. It is sobering to realize that the only time that actual meat tenderness is known is when it is eaten. We must, therefore, collectively concentrate our efforts in developing the necessary methodology to predict meat tenderness prior to eating. We are placing special emphasis on knowledge acquisition to develop such technology.

References

- ASGHAR A., BHATTI A.R. 1987. Endogenous proteolytic enzymes in skeletal muscle: Their significance in muscle physiology and during postmortem aging events in carcasses. *Adv. Food Res.*, 31, 343-451.
- BARRETT A.J., 1973. Human cathepsin B1. Purification and some properties of the enzyme. *Biochem. J.*, 131, 809-822.
- BUSCH W.A., STROMER M.H., GOLL D.E., SUZUKI A., 1972. Ca^{2+} -specific removal of Z lines from rabbit skeletal muscle. *J. Cell Biol.*, 52, 367-381.
- CROUSE J.D., CUNDIFF L.V., KOCH R.M., KOOHMARAIE M., SEIDEMAN S.C., 1989. Comparisons of Bos indicus and Bos taurus inheritance for carcass beef characteristics and meat palatability. *J. Anim. Sci.*, 67, 2661-2668.
- CROUSE J.D., SEIDEMAN S.C., CUNDIFF L.V., 1987. The effect of carcass electrical stimulation on meat obtained from Bos indicus and Bos taurus cattle. *J. Food Qual.*, 10, 407-416.
- DAVEY C.L., 1983. Post-mortem chemical changes in muscle-meat aging. *Proc. Recip. Meat Conf.*, 36, 108-115.
- DAVEY C.L., GILBERT K.V., 1969. Studies in meat tenderness. 7. Changes in the fine structure of meat during aging. *J. Food Sci.*, 34, 69-74.
- DRANSFIELD E., 1992. Modelling post-mortem tenderisation--III: Role of calpain I in conditioning. *Meat Sci.*, 31, 85-94.
- DUTSON T.R., 1983. Relationship of pH and temperature to distribution of specific muscle proteins and activity of lysosomal proteinases. *J. Food Biochem.*, 7, 223-245.
- DUTSON T.R., PEARSON A.M., 1985. Postmortem conditioning of meat. In "Advances in meat research" Vol. 1 (A.M. Pearson and T.R. Dutson, eds). AVI Publishing Co., Inc., Westport, CT, 45-72pp.
- FIEMS L.O., BUTS B., COTTYN B.G. 1990. Effect of a β -agonist on meat quality and myofibrillar protein fragmentation in bulls. *Meat Sci.*, 27, 29-39.
- GOLL D.E., OTSUKA Y., NAGAINIS P.A., SHANNON J.D. SATHE S.K., MUGURUMA, M., 1983. Role of muscle proteinases in maintenance of muscle integrity and mass. *J. Food Biochem.*, 7, 137-177.
- GREASER M.L., 1986. Conversion of muscle to meat. In "Muscle as food" (P.J. Bechtel, ed). Academic Press, NY, 37-102pp.
- HOAGLAND R., MCBRYDE C.N., POWICK W.C., 1917. Changes in fresh beef during cold storage above freezing. Bulletin No. 433, U.S. Dept. of Agric. 1-100pp.
- HOROWITS R., KEMPNER E.S., BISHOP M.E., PODOLSKY R.J., 1986. A physiological role for titin and nebulin in skeletal muscle. *Nature*, 323, 160-164.
- KOCH R.M., DIKEMAN M.E., CROUSE J.D., 1982. Characterization of biological types of cattle (Cycle III). III. Carcass composition, quality and palatability. *J. Anim. Sci.*, 54, 35-45.
- KOOHMARAIE M., 1988. The role of endogenous proteases in meat tenderness. *Proc. Recip. Meat Conf.*, 41, 89-100.

- KOOHMARAIE M., 1990. Inhibition of postmortem tenderization in ovine carcasses through infusion of zinc. *J. Anim. Sci.*, 68, 1476-1483.
- KOOHMARAIE M., 1992a. The role of Ca²⁺-dependent proteases (calpains) in postmortem proteolysis and meat tenderness. *Biochimie*, (In press).
- KOOHMARAIE M., 1992b. Role of the neutral proteinases in postmortem muscle protein degradation and meat tenderness. *Recip. Meat Conf. Proc.* 44, (In press).
- KOOHMARAIE M., 1992c. Effect of pH, temperature and inhibitors on autolysis and catalytic activity of bovine skeletal muscle μ -calpain. *J. Anim. Sci.*, (Accepted).
- KOOHMARAIE M., 1992d. Ovine skeletal muscle multicatalytic proteinase complex (proteasome): purification, characterization and comparison of its effects on myofibrils with μ -calpains. *J. Anim. Sci.*, (Submitted).
- KOOHMARAIE M., BABIKER A.S., MERKEL R.A., DUTSON T.R., 1988a. Role of Ca-dependent proteases and lysosomal enzymes in postmortem changes in bovine skeletal muscle. *J. Food Sci.*, 53, 1253-1257.
- KOOHMARAIE M., BABIKER A.S., MERKEL R.A., DUTSON T.R., 1988b. Acceleration of postmortem tenderization in ovine carcasses through activation of Ca-dependent proteases. *J. Food Sci.*, 53, 1638-1641.
- KOOHMARAIE M., CROUSE J.D., MERSMANN H.J., 1989. Acceleration of postmortem tenderization in ovine carcasses through infusion of calcium chloride. *J. Anim. Sci.*, 67, 934-942.
- KOOHMARAIE M., SHACKELFORD S.D., 1991. Effect of calcium chloride infusion on the tenderness of lambs fed a β -adrenergic agonist. *J. Anim. Sci.*, 69, 2463-2471.
- KOOHMARAIE M., SHACKELFORD S.D., MUGGLI-COCKETT N.E., STONE R.T., 1991a. Effect of the β -adrenergic agonist L644,969 on muscle growth, endogenous proteinase activities, and postmortem proteolysis in wether lambs. *J. Anim. Sci.*, 69, 4823-4835.
- KOOHMARAIE M., WHIPPLE G., CROUSE J.D., 1990. Acceleration of postmortem tenderization in lamb and Brahman-cross beef carcasses through infusion of calcium chloride. *J. Anim. Sci.*, 68, 1278-1283.
- KOOHMARAIE M., WHIPPLE G., KRETCHMAR D.H., CROUSE J.D., MERSMANN H.J., 1991b. Postmortem proteolysis in longissimus muscle from beef, lamb and pork carcasses. *J. Anim. Sci.*, 69, 617-624.
- KRETCHMAR D.H., HATHAWAY M.R., DAYTON W.R., 1990. Alterations in postmortem degradation of myofibrillar proteins in muscle of lambs fed a β -adrenergic agonist. *J. Anim. Sci.*, 68, 1760-1772.
- KUMAMOTO T., KLEESE W.C., CONG J.Y., GOLL D.E., 1992. Localization in electron micrographs of the Ca-dependent proteinases and their inhibitor in normal, starved, and denervated rat skeletal muscle. *Anat. Rec.*, 232, 60-77.
- LaCOURT A.A., OBLED A., DEVAL C., OUALI A., VALIN C., 1986. Postmortem localization of lysosomal peptide hydrolase. Cathepsin B. In "Cysteine proteinases and their inhibitors" (Walter de Gruyter and Co., eds). New York, 239-248pp.
- LEHMANN K.B., 1907. Studies of the causes for the toughness of meats. *Arch. Hyg.*, 3, 134.
- MARSH B.B., 1977. The nature of tenderness. *Recip. Meat Conf. Proc.*, 30, 69-74pp.
- MARSH B.B., 1983. Effects of early-postmortem muscle pH and temperature on meat tenderness. *Recip. Meat Conf. Proc.*, 36, 131-135pp.
- MARSH B.B., RINGKOB, T.P., RUSSELL, R.L., SWARTZ, D.R., PAGEL, L.A., 1988. Mechanisms and strategies for improving meat tenderness. *Recip. Meat Conf. Proc.* 41, 113-118pp.
- MORGAN J.B., MILLER R.K., MENDEZ F.M., HALE D.S., SAVELL J.W., 1991. Using calcium chloride injection to improve tenderness of beef from mature cows. *J. Anim. Sci.*, 69, 4469-4476.
- ORLOWSKI M., 1990. The multicatalytic proteinase complex, a major extralysosomal proteolytic system. *Biochemistry*, 29, 10289-10297.
- OUALI A., 1990. Meat tenderization: Possible causes and mechanisms. A Review. *J. Musc. Foods.*, 1, 129-165
- OUALI A., 1992. Proteolytic and physicochemical mechanisms involved in meat texture development. *Biochimie*, (In press).
- PARRISH F.C., JR., 1977. Skeletal muscle tissue disruption. *Recip. Meat Conf. Proc.* 36, 87-98pp.
- PEACOCK F.M., KOGER M., PALMER A.Z., CARPENTER J.W., OLSON T.A., 1982. Additive breed and heterosis effects for individual and maternal influences on feedlot gain and carcass traits of Angus, Brahman, Charolais and crossbred steers. *J. Anim. Sci.*, 55, 797-803.

- PEARSON A.M., 1986. Physical and biochemical changes occurring in muscle during storage and preservation. In "Muscle as food" (P.J. Bechtel, ed). Academic Press, NY, 103-133pp.
- PENNY I.F., 1980. The enzymology of conditioning. In "Developments in Meat Science-1" (R. Lawrie, ed). Applied Science Publishers, Ltd., London, 115-143pp.
- PRINGLE T.D., CALKINS C.R. KOOHMARAIE M., JONES S.J., 1992. Response time of animal performance, muscle growth, endogenous muscle proteinase activities, muscle metabolism and meat tenderness to β -adrenergic agonist feeding in wether lambs. J. Anim. Sci., (Submitted).
- RAMSEY C.B., COLE J.W., MEYER B.H., TEMPLE R.S., 1961. Effects of type and breed of British, Zebu and dairy cattle on production, palatability and composition. II. Palatability differences and cooking losses as determined by laboratory and family panels. J. Anim. Sci., 22, 1001-1008.
- RIVETT A.J., 1989. The multicatalytic proteinase of mammalian cells. Arch. Biochem. Biophys., 268, 1-8.
- ROBSON R.M., HUIATT, T.W. 1983. Roles of the cytoskeletal proteins desmin, titin and nebulin in muscle. Recip. Meat Conf. Proc. 36, 116-124pp.
- ROBSON R.M., O'SHEA J.M., HARTZER M.K., RATHBUN W.E., LaSALLE F., SCHREINER P.J., KASANG L.E., STROMER M.H., LUSBY M.L., RIDPATH J.F., PANG Y.Y., EVANS R.R., ZEECE M.G., PARRISH F.C., HUIATT T.W. 1984. Role of new cytoskeletal elements in maintenance of muscle integrity. J. Food Biochem. 8, 1-24pp.
- ROBSON R.M., YAMAGUCHI, M., HUIATT, T.W., RICHARDSON, F.L., O'SHEA, J.M., HARTZER, M.K., RATHBUN, W.E., SCHREINER, P.J., STROMER, M.H., PANG, S., EVANS, R.R., RIDPATH, J.F. 1981. Biochemistry and molecular architecture of muscle cell 10-nm filaments and Z-line. Roles of desmin and alpha-actinin. Recip. Meat Conf. Proc. 34, 5-11pp.
- SHACKELFORD S.D., KOOHMARAIE M., CROUSE J.D., MILLER M.F., REAGAN J.O., 1991. An evaluation of tenderness of the longissimus muscle of Angus by Hereford versus Brahman crossbred heifers. J. Anim. Sci., 69, 171-177.
- ST. ANGELO A.J., KOOHMARAIE M. CRIPPEN K.L., CROUSE J.D., 1991. Acceleration on tenderization/inhibition of warmed-over flavor by calcium chloride-antioxidant infusion into lamb carcasses. J. Food Sci., 56, 359-362.
- TARRANT P.V., 1987. Review: Muscle Biology and Biochemistry. Proc. Int. Cong. Meat Sci. Technol., 33, 1-5.
- TAYLOR M.A.J., ETHERINGTON D.J., 1991. The solubilization of myofibrillar proteins by calcium ions. Meat Sci., 29, 211-219.
- WANG, K., 1985. Sarcomere-associated cytoskeletal lattices in striated muscle. Review and hypothesis. Cell Muscle Motil., 6, 315-369.
- WANG K., McCARTER R., WRIGHT J., BEVERLY J., RAMIREZ-MITCHELL R., 1991. Regulation of skeletal muscle stiffness and elasticity by titin isoforms: A test of the segmental extension model of resting tension. Proc. Natl. Acad. Sci. (USA), 88, 7101-7105.
- WHEELER T.L., CROUSE, J.D., KOOHMARAIE M., 1992. The effect of postmortem injection time and freezing on the effectiveness of calcium chloride at improving meat tenderness. J. Anim. Sci., (Submitted).
- WHEELER T.L., KOOHMARAIE M., 1992. Effects of the β -adrenergic agonist L644,969 on muscle protein turnover, endogenous proteinase activities and meat tenderness in steers. J. Anim. Sci., (Accepted).
- WHEELER T.L., KOOHMARAIE M., CROUSE J.D., 1991. Effects of calcium chloride injection and hot-boning on the tenderness of round muscles. J. Anim. Sci., 69, 4871-4875.
- WHEELER T.L., SAVELL J.W., CROSS, H.R., LUNT D.K., SMITH S.B., 1990. Mechanisms associated with the variation in tenderness of meat from Brahman and Hereford cattle. J. Anim. Sci., 68, 4206-4220.
- WHIPPLE G., KOOHMARAIE M., 1992. Calcium chloride in vitro effects on isolated myofibrillar proteins. J. Anim. Sci., (Submitted).
- WHIPPLE, G., KOOHMARAIE, M., DIKEMAN, M.E., CROUSE J.D., HUNT M.C., KLEMM R.D., 1990. Evaluation of attributes that affect longissimus muscle tenderness in Bos taurus and Bos indicus cattle. J. Anim. Sci., 68, 2716-2728.
- WINGER R.J., POPE C.G., 1980-81. Osmotic properties of post-rigor beef muscle. Meat Sci., 5, 355-369.
- YOSHIOKA T., HIGUCHI H., KIMURA S., OHASHI K., UMAYAMA Y., MARUYAMA K., 1986. Effects of mild trypsin treatment on the passive tension generation and connectin splitting in stretched skinned fibers from frog rana-catesbeiana skeletal muscle. Biomed. Res. 7, 181-186.