Table 1. Breakpoints Used for Susceptibility Testing of *Campylobacter* [1]

CLSI Subclass[2]	Antimicrobial Agent	Antimicrobial Concentration Range (µg/ml) 1998-2004	Breakpoints (µg/ml) Etest (1998-2004)			Antimicrobial Concentration Range (µg/ml) 2005-2012	Breakpoints (µg/ml) Microbroth Dilution (2005-2012)		
			Susceptible	Intermediate	Resistant		Susceptible	Intermediate	Resistant
Aminoglycosides	Gentamicin	0.016 - 256	≤ 4	8	≥ 16	0.12 - 32	≤ 2	4	≥ 8
Lincosamides	Clindamycin	0.016 - 256	≤ 0.5	1 - 2	≥ 4	0.03 - 16	≤ 2	4	≥ 8
Macrolides	Azithromycin	0.016 - 256	≤ 0.25	0.5 - 1	≥ 2	0.015 - 64	≤ 2	4	≥ 8
	Erythromycin	0.016 - 256	≤ 0.5	1 - 4	≥ 8	0.03 - 64	≤ 8	16	≥ 32
Ketolides	Telithromycin	NT	NT	NT	NT	0.015 - 8	≤ 4	8	≥ 16
Phenicols	Florfenicol	NT	NT	NT	NT	0.03 - 64	≤ 4	N/A	N/A
	Chloramphenicol	0.016 - 256	≤ 8	16	≥ 32	NT	NT	NT	NT
Fluoroquinolones	Ciprofloxacin	0.002 - 32	≤ 1	2	≥ 4	0.015 - 64	≤ 1	2	≥ 4
Quinolones	Nalidixic acid	0.016 - 256	≤ 16	N/A	≥ 32	4 - 64	≤ 16	32	≥ 64
Tetracyclines	Tetracycline	0.016 - 256	≤ 4	8	≥ 16	0.06 - 64	≤ 4	8	≥ 16

[1] Breakpoints established by CLSI (Clinical and Laboratory Standards Institute) were used when available. CLSI breakpoints are available only for erythromycin, ciprofloxacin, and tetracycline

[2] According to CLSI M100 document

N/A - Not applicable

NT- Not tested